



Lake County Substance Use and Consequences, 2011

MN STATE EPIDEMIOLOGICAL OUTCOMES WORKGROUP

KEY FINDINGS

- The average cost per capita of alcohol-related motor vehicle crashes, fatalities and injuries in Lake County is more than double the state average.
- The average annual lung, bronchus and trachea cancer death rate for Lake County is higher than both the Northeast and state rates.
- The percent of Lake County narcotics arrests involving marijuana have been on the decline in the past three years.

INSIDE

Tobacco Consumption 2

Tobacco Consequences..... 2

Other Drug Consumption 3

Other Drug Consequences... 3

MN SEOW 4

Data Sources..... 4

Alcohol Consumption

Reported past-month alcohol use among students dropped in the Northeast Alcohol, Tobacco and Other Drug Prevention Region from 34% in 1998 to 25% in 2007. Comparatively, the state rate dipped from 31% to 23%. The percent of students reporting past-two week binge drinking also declined during that period—Northeast:


28% to 22%, Minnesota 26% to 20% (MSS).

Among adults, reported 30-day alcohol use was lower than the state average for Northeast men (66% vs. 67%) and women (48% vs. 53%) in 2004/2005. While women in the Northeast Region were slightly less likely to report past-month binge drinking

than the state average, 12% vs. 13%, Northeast men were more likely to do so: 27% vs. 24% (MNSASU).

Minnesota Student Survey data on use of alcohol by Lake County students are not available for 2001 to 2010. Statistics on adult alcohol consumption are not available at the county level.

Alcohol Consequences



From 2005 to 2009, the average cost per capita of alcohol-related motor vehicle crashes, fatalities and injuries was \$127 in Lake County, compared to \$96 in the Northwest and \$54 in the state (OTS).


Over that same period, the driving while intoxicated (DWI) arrest rate with Lake as

the county of residence averaged 62.0 per 10,000 population. With Lake as the county of arrest, the average rate was 57.0 (OTS). For comparison, the state DWI rate averaged 71.3 per 10,000.

The percent of all motor vehicle crashes that were alcohol-related averaged 6%

in Lake County, 8% in the Northeast Region, and 5% in the state (OTS).

The cirrhosis death rate per 100,000 population in Lake County averaged 9 from 2004 to 2009, as compared to 12 in the Northeast and 6 in the state (MCHS).



The US Census Bureau estimate for the 2008 Lake County population is 10,609. Lake County is approximately 98% White, 1% American Indian/Alaska Native and 1% Hispanic/Latino. In 2000, the percent of individuals living below the poverty level was 7% in Lake County, as compared to 11% in the Northeast Region and 8% in Minnesota.

Tobacco Consumption

Reported 30-day cigarette smoking among Northeast students, though on the decline, has been higher than the state average over the past 9 years. Northeast rates dropped for: 9th grade males, 25% to 15%; 9th grade females, 29% to 16%; 12th grade males, 38% to 27%; and 12th grade females, 40% to 27% (MSS).

2004/2005 reported past-month smoking rates were higher than the state average for adults in the Northeast: 26% vs. 23%. However, Northeast rates were lower

than average for 18–24 year-old men and women aged 21–24 (MNSASU).

The percent of male students reporting past-month chewing tobacco use in 2007 was higher in the Northeast Region than the state average for 6th graders, 2% vs. 1%; 9th graders, 15% vs. 8%; and 12th graders, 25% vs. 19% (MSS).


Northeast students were slightly more likely than the state average to report first having smoked all or part of a cigarette by age 13 or younger:

17% vs. 16% for males and 16% vs. 14% for females.

Similarly, Northeast male students were more likely to report first having smoked a cigar or used chewing tobacco by the age of 13 or younger: 14% vs. 9% (MSS).

Minnesota Student Survey data on use of tobacco by Lake County students are not available for 2001 to 2010. Statistics on adult alcohol consumption are not available at the county level.

Tobacco Consequences



The average annual rate per 100,000 population of lung, bronchus and trachea cancer deaths in Lake County was 80 from 2004 to 2008. Comparatively, the Northeast averaged 69 per 100,000 and Minnesota averaged 45 per 100,000 (MCHS).

The number of Lake County lung and bronchus cancer deaths averaged 4 per year among males and females from 2004 to 2008 (MCSS). Note: these rates exclude mesothelioma (ICD 10 Codes C45).

It is estimated that 90% of lung cancer deaths among males and 79% of lung cancer deaths among females in the United States are smoking-related.

Other Drug Consumption

The percent of Northeast students reporting past-month marijuana use dropped from 15% in 1998 to 11% in 2007, and state rates from 12% to 9%. From 2001 to 2007, reported rates of methamphetamine use dipped from 8% to 2%; MDMA/Ecstasy use from 5% to 3%; and crack/cocaine use from 7% to 4% in the Northeast Region (MSS).

In 2007, 9th and 12th grad-

ers were asked about abuse of specific types of prescription drugs. Northeast students reported higher than average rates of abuse: painkillers, 6% vs. 5%, ADD/ADHD drugs, 6% vs. 4%, and stimulants/diet pills: 4% vs. 2% (MSS).


Reported past-month use of marijuana was slightly lower than the state average among Northeast adults: 3% vs. 4%, as was reported abuse of MDMA/

Ecstasy: 0% vs. 1%. Reported past-year use among Northeast adults equaled the state rate for pain relievers, 2%, crack/cocaine, 1%, and methamphetamines, 1% (MNSASU).

Minnesota Student Survey data on use of drugs by Lake County students are not available for 2001 to 2010. Statistics on adult alcohol consumption are not available at the county level.


The number of Lake County males admitted to Minnesota treatment facilities increased from 37 in 2000 to 44 in 2008; the number of females admitted increased from 14 to 16. In 2008, 10% of Lake County residents admitted to treatment programs were under age 18; 17% were between the ages of 18 and 24; 42% were aged 25-44 and 32% were aged 45-64.

After alcohol, the second most common primary substance of abuse for Lake County residents was marijuana.



Other Drug Consequences

There were 14 Lake County narcotics arrests in 2009—79% of which involved marijuana (BCA). From 2007 to 2009, the average annual rate of adults on probation for drug offenses in Lake County was 26.0 per 10,000 population; there were no juveniles reported to be on probation for drug offenses. The rate of adults in prison for drug offenses averaged 2.7 per 10,000 population during that same period (DOC).



MN STATE EPIDEMIOLOGICAL OUTCOMES WORKGROUP

Questions or comments? Contact:

Melissa Boeke, MS
Epidemiologist

Minnesota Institute of Public Health
2720 Highway 10 NE
Mounds View, MN 55112

Phone: 763.712.7615
Fax: 763.427.7841
E-mail: sumn@miph.org

What's Happening in Your County?



Substance Use in Minnesota, located at www.sumn.org, puts data on alcohol, tobacco and other drug use and consequences at your fingertips. The Web site was designed to help communities make decisions about substance abuse prevention efforts based on 55 indicators.

This site was developed by the Minnesota State Epidemiological Outcomes Workgroup (SEOW) with funding from the Minnesota Department of Human Services, Alcohol and Drug Abuse Division (ADAD). SEOW membership includes: Minnesota's departments of Human Services, Health, Education, Public Safety and Corrections and the Minnesota Institute of Public Health, which maintains the Web site.

Data Sources

Population statistics are from the U.S. Census Bureau.

Youth alcohol, tobacco and other drug consumption data are from the Minnesota Student Survey (MSS). The MSS is a confidential and anonymous self-administered survey given every three years to 6th, 9th and 12th grade students attending Minnesota public, charter and tribal schools. The Minnesota Student Survey Interagency Team is made up of four state agencies: the Minnesota Department of Education, the Minnesota Department of Health, the Minnesota Department of Human Services, and the Minnesota Department of Public Safety. Adult consumption data are from the MN Survey of Adult Sub-

stance Use (MNSASU) from the Minnesota Department of Human Services, Performance Measurement and Quality Improvement. Binge drinking is defined as 5 or more drinks in a row on one occasion in the MSS, and 5 or more drinks for males or 4 or more drinks for females in the MNSASU.

Alcohol consequence data are from the Office of Traffic Safety's (OTS) *Impaired Driving Facts and Crash Facts*, and from the National Safety Council.

Alcohol and tobacco mortality data are from the Minnesota Center for Health Statistics (MCHS) and the Minnesota Cancer Surveillance System (MCSS).

Drug consequence data are

from the Minnesota Department of Corrections' (DOC) *Inmate Profile and Probation Survey*, and from the Bureau of Criminal Apprehension's (BCA) *Minnesota Crime Information* reports.

Treatment data are from the Drug and Alcohol Abuse Normative Evaluation System (DAANES).

Detailed data source information can be found at www.sumn.org. Data source fact sheets can be found under Tools. Links to each data source, when available, can be found under each table when using Data by Location or Data by Topic.

Aggregated data at the state and county level do not reveal disparities that may exist within a given geographic area.