



Martin County Substance Use and Consequences, 2011

MN STATE EPIDEMIOLOGICAL OUTCOMES WORKGROUP

KEY FINDINGS

- Martin County's average cost per capita of alcohol-related motor vehicle crashes, fatalities and injuries is higher than both the regional and state averages.
- From 1998 to 2010, youth smoking rates fell from 23% to 9% in Martin County.
- Reported past-year use of methamphetamines dropped among Martin students from 7% in 2004 to 2% in 2010.

INSIDE

Tobacco Consumption 2
 Tobacco Consequences..... 2
 Risk & Protective Factors 2
 Other Drug Consumption 3
 Other Drug Consequences... 3
 MN SEOW 4
 Data Sources..... 4

Alcohol Consumption

The rate of reported past-month alcohol use by youth in Martin County dropped from 36% in 1998 to 17% in 2010. Comparatively, the rate for Minnesota's Southwest Alcohol, Tobacco and Other Drug Prevention Region fell from 33% to 21%, and the states from 31% to 20%.

Reported use varied by grade (*see graph below*). In


2010, Martin County female students were less likely than male students to report past month use: 14% vs. 19%.

The percent of Martin County students reporting binge drinking in the past two weeks—having five or more drinks in a row on one occasion—fell from 30% in 1998 to 11% in 2010. In comparison, the Southwest rate fell

from 28% to 17% and the state rate from 26% to 16%.

In 2010, 7% of Martin County 9th graders reported binge drinking, as did 15% of 12th graders. Binge drinking was reported by more male students than female students: 12% vs. 10% (MSS).

Statistics on adult alcohol consumption are not available at the county level.



Alcohol Consequences

From 2005 to 2009, the average cost per capita of alcohol-related motor vehicle crashes, fatalities and injuries was \$85 in Martin County, \$70 in the Southwest Region and \$54 in Minnesota (OTS).


Over that same period, the driving while intoxicated

(DWI) arrest rate with Martin as the county of residence averaged 65.5 per 10,000 population. With Martin as the county of arrest, the average rate was 68.4 (OTS).

The percent of all motor vehicle crashes that were alcohol-related averaged 6% in

Martin County and the Southwest Region, compared to 5% in Minnesota (OTS).

The annual cirrhosis death rate per 100,000 population averaged 6 in both Martin County and in the state, and 7 in the Southwest Region from 2004 to 2008 (MCHS).



Statistics on adult substance use are available by region for 2004/2005:

- Past-month alcohol use was lower than the state average for Southwest women, 50% vs. 53%, but equal for men: 67%.

- In the Southwest, past month smoking was reported by 41% of 18-20 year-olds, 41% of 21-24 year-olds, 26% of 25-44 year-olds, 19% of 45-64 year-olds, and 8% of persons aged 65+.

- Southwest adults were less likely than the state average to report past month use of marijuana, 2% vs. 4%. They were as likely as the state average to report past year use of crack/cocaine, 1%, but less likely to report past-year use abuse of prescription pain relievers: 1% vs. 2%. Less than one percent of Southwest adults reported past-year methamphetamine use.

Minnesota Survey of Adult Substance Use

Tobacco Consumption

Smoking among youth declined statewide from 1998 to 2010: down from 23% to 9% in Martin County, 20% to 12% in the Southwest Region, and 19% to 10% in Minnesota.

In 2010, Martin County rates varied greatly by grade (*see graph*). Past month smoking was reported by 8% of Martin male students and 11% of female students in 2010.


From 2004 to 2010, reported past-month chewing tobacco use increased among

Martin County males: 4% to 10%.

Martin youth were a bit more likely than average to report first smoking by age 13

or younger: 13% vs. 12% (MSS).

Statistics on adult smoking are not available at the county level.



Tobacco Consequences

The average annual rate per 100,000 population of lung, bronchus and trachea cancer deaths in Martin County was 60 from 2004 to 2008. Comparatively, the Southwest rate averaged 51 and the state rate

averaged 45 (MCHS).

The number of Martin County lung and bronchus cancer deaths averaged seven among males and five among females from 2004 to 2008 (MCSS).

It is estimated that 90% of lung cancer deaths among males and 79% of lung cancer deaths among females in the United States are smoking related.

Risk and Protective Factors

In 2010, students reporting past 30 day use of alcohol and tobacco were asked how they obtained those substances. Martin County students most often reported social access to alcohol (82%), and most often reported buying (62%) or getting (56%) tobacco from someone or somewhere. Martin County students were less likely (17%) than Southwest (23%) and Minnesota students (26%) to report taking alcohol

from someone or somewhere, and a bit less likely to report taking tobacco (13%, 14%, 17%) from someone or somewhere.

Students were also asked how much they thought people risked harming themselves by frequently binge drinking, smoking one or more packs of cigarettes per day, or smoking marijuana once or twice per week. Perception of great or moderate risk of alcohol harm was reported by 80% of Martin

County students, perception of tobacco harm was reported by 89% of Martin County students, and perception of marijuana harm was reported by 80% of Martin students. Similarly, students were asked if they thought their parents or guardians would disapprove or greatly disapprove of such levels of substance use. Perception of disapproval was 88% for alcohol, 94% for tobacco and 97% for marijuana (MSS).

Other Drug Consumption

Martin County students are more likely to report use of marijuana than any other illicit drug (*see graph below*). In 2010, marijuana use was reported by 1% of Martin County 6th graders, 4% of 9th graders and 11% of 12th graders. Male students were less likely than female students to report use: 4% vs. 6%.

From 2004 to 2010, reported past-year use of methamphetamines fell from 5% to 4% among Martin 9th graders, and 9% to 0% among 12th graders. From 1998 to 2010,


reported past-year use of inhalants fell among Martin 6th graders, 7% to 6%, 9th graders, 9% to 5%, and 12th graders, 4% to 1%. During that period, reported past-year use of psychedelics fell among male students, 13% to 2%, holding for females at 3%. Reported use of crack/cocaine fell from 10% to 1% for males and held steady at 3% for females.

In 2010, 2% of Martin 6th graders reported past-year abuse of prescription drugs compared to 1% in the Southwest Region and in the state.

Martin County 9th and 12th graders were asked about abuse of specific types of prescription drugs in 2010: painkillers (reported by 5% of 9th graders and 3% of 12th graders), ADD/ADHD drugs (4% of 9th graders; 2% of 12th graders), tranquilizers/sedatives (4% of 9th graders; 2% of 12th graders), and stimulants/diet pills: 4% of 9th graders; 1% of 12th graders (MSS).

Statistics on adult drug use are not available at the county level.

The US Census Bureau estimate for the 2008 Martin County population is 21,780. Martin County is approximately 97% White, 2% Hispanic/Latino. In 2000, the percent of individuals living below the poverty level was 11% in Martin County compared to 9% in the Southwest Region and 8% in the state.



sequences

There were Martin County narcotics arrests in 74 in 2009—42 of which involved marijuana. In comparison, the average annual number of narcotics arrests for Martin County from 2005 to 2009 was 84 (BCA). For probation and prison data related to drug offenses, see table below (DOC).

DRUG OFFENSES	MARTIN COUNTY			SOUTHWEST REGION			MINNESOTA		
Rates per 1,000 Population	2007	2008	2009	2007	2008	2009	2007	2008	2009
Adults on Probation	4.4	5.0	4.9	2.6	4.2	3.1	3.4	3.6	3.5
Juveniles on Probation	0.7	1.4	0.4	0.3	0.2	0.6	0.4	0.5	0.5
Adults in Prison	1.0	1.0	0.6	0.5	0.6	0.4	0.4	0.5	0.3

MN STATE EPIDEMIOLOGICAL OUTCOMES WORKGROUP

Questions or comments? Contact:

Melissa Boeke, MS
Epidemiologist

Minnesota Institute of Public Health
2720 Highway 10 NE
Mounds View, MN 55112

Phone: 763.712.7615

Fax: 763.427.7841

E-mail: sumn@miph.org

What's Happening in Your County?



Substance Use in Minnesota, located at www.sumn.org, puts data on alcohol, tobacco and other drug use and consequences at your fingertips. The Web site was designed to help communities make decisions about substance abuse prevention efforts based on 55 indicators.

This site was developed by the Minnesota State Epidemiological Outcomes Workgroup (SEOW) with funding from the Minnesota Department of Human Services, Alcohol and Drug Abuse Division (ADAD). SEOW membership includes: Minnesota's departments of Human Services, Health, Education, Public Safety and Corrections and the Minnesota Institute of Public Health, which maintains the Web site.

Data Sources

Population statistics are from the U.S. Census Bureau.

Youth alcohol, tobacco and other drug consumption data are from the Minnesota Student Survey (MSS). The MSS is a confidential and anonymous self-administered survey given every three years to 6th, 9th and 12th grade students attending Minnesota public, charter and tribal schools. The Minnesota Student Survey Interagency Team is made up of four state agencies: the Minnesota Department of Education, the Minnesota Department of Health, the Minnesota Department of Human Services, and the Minnesota Department of Public Safety. Adult consumption data are

from the MN Survey of Adult Substance Use (MNSASU) from the Minnesota Department of Human Services, Performance Measurement and Quality Improvement. Binge drinking is defined as 5 or more drinks in a row on one occasion in the MSS, and 5 or more drinks for males or 4 or more drinks for females in the MNSASU.

Alcohol consequence data are from the Office of Traffic Safety's (OTS) *Impaired Driving Facts and Crash Facts*, and from the National Safety Council.

Alcohol and tobacco mortality data are from the Minnesota Center for Health Statistics (MCHS) and the Minnesota Cancer Surveillance

System (MCSS).

Drug consequence data are from the Minnesota Department of Corrections' (DOC) *Inmate Profile and Probation Survey*, and from the Bureau of Criminal Apprehension's (BCA) *Minnesota Crime Information* reports.

Detailed data source information can be found at www.sumn.org. Data source fact sheets can be found under Tools. Links to each data source, when available, can be found under each table when using Data by Location or Data by Topic.

Aggregated data at the state and county level do not reveal disparities that may exist within a given geographic area.